

REVIEW

Application of a new system for classifying root and canal anatomy in studies involving micro-computed tomography and cone beam computed tomography: Explanation and elaboration

H. M. A. Ahmed¹ , N. Ibrahim², N. S. Mohamad³, P. Nambiar^{2,4}, R. F. Muhammad⁵, M. Yusoff⁵ & P. M. H. Dummer⁶ 

¹Department of Restorative Dentistry, Faculty of Dentistry, University of Malaya, Kuala Lumpur; ²Department of Oral and Maxillofacial Clinical Sciences, Faculty of Dentistry, University of Malaya, Kuala Lumpur; ³Centre of Medical Imaging, Faculty of Health Sciences, Universiti Teknologi MARA (UiTM) Selangor Branch, Puncak Alam Campus, Bandar Puncak Alam, Selangor; ⁴Department of Oral Biology and Biomedical Sciences, Faculty of Dentistry, MAHSA University, Bandar Saujana Putra; ⁵Department of Geology Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia; and ⁶School of Dentistry, College of Biomedical and Life Sciences, Cardiff University, Cardiff, UK

Abstract

Ahmed HMA, Ibrahim N, Mohamad NS, Nambiar P, Muhammad RF, Yusoff M, Dummer PMH. Application of a new system for classifying root and canal anatomy in studies involving micro-computed tomography and cone beam computed tomography: Explanation and elaboration. *International Endodontic Journal*, **54**, 1056–1082, 2021.

Adequate knowledge and accurate characterization of root and canal anatomy is an essential prerequisite for successful root canal treatment and endodontic surgery. Over the years, an ever-increasing body of knowledge related to root and canal anatomy of the human dentition has accumulated. To correct deficiencies in existing systems, a new coding system for classifying root and canal morphology, accessory canals and anomalies has been introduced. In recent

years, micro-computed tomography (micro-CT) and cone beam computed tomography (CBCT) have been used extensively to study the details of root and canal anatomy in extracted teeth and within clinical settings. This review aims to discuss the application of the new coding system in studies using micro-CT and CBCT, provide a detailed guide for appropriate characterization of root and canal anatomy and to discuss several controversial issues that may appear as potential limitations for proper characterization of roots and canals.

Keywords: accessory canals, anatomy, anomalies, canal configuration, cone beam computed tomography, micro-computed tomography, new system, root, root canal morphology.

Received 2 December 2020; accepted 27 January 2021

Introduction

Root and canal morphology have been the subject of numerous studies using a wide variety of techniques including ground extracted teeth, staining and clearing extracted teeth, conventional 2D radiographic

images and the more recently 3D imaging techniques including micro-computed tomography (micro-CT) and cone beam computed tomography (CBCT) (Weine *et al.* 1969, 1999, Vertucci 1984, Versiani *et al.* 2016, Martins *et al.* 2017, Ahmed & Ross-Fedele 2020). The application of micro-CT in laboratory-

Correspondence: Hany Mohamed Aly Ahmed, Department of Restorative Dentistry, Faculty of Dentistry, University of Malaya, 50603 Kuala Lumpur, Malaysia (e-mail: hany_endodontist@hotmail.com).

based experimental studies has generated considerable and more detailed information (compared to other methods) on root and canal morphological characteristics such as accessory roots (Versiani *et al.* 2012), root canal configurations (Leoni *et al.* 2014, Filpo-Perez *et al.* 2015), canal isthmuses (Keleş & Keskin 2018), accessory canals (Xu *et al.* 2016, 2019) and root canal anomalies (Gu 2011, Zhang *et al.* 2014). In clinical settings, high resolution CBCT has been used to study root and canal morphology in various population groups (Abella *et al.* 2015, Martins *et al.* 2017, Kantilieraki *et al.* 2019, Karobari *et al.* 2020), as well as being a useful tool for detection of missed canals, and managing cases with complex canal morphology (Kotthoor *et al.* 2010, Badole *et al.* 2014, Hashem & Ahmed 2017).

A new coding system for classifying the root and canal anatomy has been proposed which provides detailed information on tooth notation, number of roots and root canal configuration in addition to accessory canals and dental anomalies (Ahmed *et al.* 2017, 2018, Ahmed & Dummer 2018a,b). This coding system overcomes deficiencies in previous systems by addressing the number of roots in every tooth type and has the ability to describe root canal configurations without referring to specific Roman numerals, which is a challenge when applying the Vertucci classification to teeth with complex canal systems (Ahmed *et al.* 2017, Saber *et al.* 2019).

Recent surveys amongst dental students and dental practitioners have supported the application of the new coding system in teaching, research and clinical practice (Ahmed *et al.* 2020a, Salas *et al.* 2021). This review aims to provide a detailed description of the application of the new coding system when undertaking laboratory-based micro-CT studies and *in vivo* studies using CBCT with an emphasis on the basic concepts to be followed. The review also addresses several controversies which remain over root, canal and pulp chamber anatomy and how they can be categorized accurately using the new coding system.

Application of the new coding system in micro-CT studies

The number of roots and their morphological variations

The inability to define the number of roots in anterior and premolar teeth together with the fact that many root canals cannot be classified are the main




Figure 1 Reconstructed micro-CT images showing the use of the new system to describe the number of roots in different teeth – the superscript on the left of the TN (Tooth Number) denotes the number of roots.

deficiencies of the Vertucci classification (Ahmed *et al.* 2017, Ahmed & Dummer 2018b, Saber *et al.* 2019). The new coding system describes the anatomical features of roots in a consistent manner regardless of the tooth type and whether a tooth is single or multi rooted (Ahmed *et al.* 2017, Ahmed & Dummer 2018b; Fig. 1).

Roots can bifurcate when the Hertwig's Epithelial Root Sheath (HERS) divides to form two similar roots, or by folding of the HERS to form an independent root which may have various morphological features (Ahmed & Abbott 2012). The new system for classifying root and canal morphology has defined that any 'division' of a root, whether in the coronal, middle or apical third, is coded as two or more roots (Ahmed *et al.* 2017; Fig. 2).

It is generally acknowledged that teeth with bifurcations in the coronal and middle thirds of roots clearly have two roots; however, this categorization is more controversial when the bifurcation is in the apical third of the root, particularly when near the root apex (Fig. 2). Turner (1981) classified bifurcations in the apical portion of the root in two forms: (i) a single-rooted tooth with bifid apex in which the bifurcation is less than one-third to one-fourth of the total root length (Fig. 2b), and (ii) a single-rooted tooth with a double apex which does not have a very clear bifurcation but has two distinct small root tips that can be identified (Fig. 2c). A recent micro-CT study on mesial roots of mandibular first molars reported that the presence of a bifid-tipped root may be associated with complex canal anatomy with a high incidence of accessory canals (Keleş *et al.* 2020; Fig. 2).




Figure 2 Micro-CT images for teeth with (a) Double roots, (b) Bifid-tipped roots (BR), (c) Double apex (DA) using the new classification system in single rooted (red box) and double rooted (blue box).

On the other hand, a micro-CT study on double-rooted mandibular canines considered bifurcations in the middle and apical thirds of the root as two separate roots (Versiani *et al.* 2011).

From the discussion above, it seems the main challenge that researchers always face is how to make the identification process simple yet accurate and consistent when an apical bifurcation of the root occurs.

The new system can be applied to describe this anatomy as follows:

1. Those roots with clear apical root bifurcations can be referred to as double rooted that usually have a common canal coronal to the level of the bifurcation (Versiani *et al.* 2011, Ahmed *et al.* 2017, Fig. 2a).
2. Those roots with apical bifurcations with no 'distinct double roots' can be categorized as either (a) single rooted with bifid tip [if the bifurcation is located in the middle portion of the apical third of the root – Bifid Root (BR)] or (b) small double apex roots [in instances of double root tips – Double Apex (DA)] as shown in Fig. 2b, c. If researchers prefer to refer to this anatomy using the new classifying system, then an abbreviation can be used as a superscript on the left of the TN if the tooth has a single root or as a superscript on the right of the TN but on the left of that specific root if the tooth is double/multi rooted (Fig. 2b, c). For simplicity, researchers could consider a root with a bifid tip or double apex as one group (single rooted) if this feature is not relevant to the specific research study.

For root fusions, the new system has complementary codes to describe teeth with roots that fuse (Fig. 3). However, it is worth noting that the literature is unclear on how fused roots are identified and classified. For instance, Hou & Tsai (1994) divided root fusion into three grades:

- Grade I:* fusion involving the cervical half of roots;
- Grade II:* fusion involving the cervical two-thirds of roots;
- Grade III:* complete or true fusion of roots.

Root fusion by cementum in the apical region only has been considered as pseudo-fusions (Hou & Tsai 1994). Any combination of grades with 1, 2, or 3 affected surfaces in maxillary molars was recorded as one-, two-, and three-surface fusions (Hou & Tsai 1994).

Ross & Evanchik (1981) defined any molar that had one root or whose roots were fused apical to the usual furcal position as a molar with fused roots. This included molars with fusion of one-third or less of the roots, and molars with fusion of entire root surfaces. Molars with roots fused only in the apical one-third and with a normal furcation were included in the category of fused teeth (Ross & Evanchik 1981).

Others have defined fusion when the ratio of the distance from the cementoamel junction (CEJ) to the apical point of root furcation or root fusion (CEJ-RF), and from the CEJ to the apex of the root (CEJ-Apex) is not less than 70% (Zhang *et al.* 2014). The authors reported that this percentage was selected based on a pilot study on maxillary molars. Even though this is considered a reasonable and reproducible approach, this percentage may vary in different teeth and population groups. The same authors also divided root fusions in maxillary molars into six groups, which is believed to be a practical approach for classifying this anatomy in three-rooted teeth (Ahmed & Dummer 2018).

This confusion in terminology reveals that the controversy over the definition of root fusions can lead to misleading conclusions. In addition, the literature reveals that root fusion is considered a normal anatomical variation by some authors, not one of the anomalies or malformations that may affect roots




Figure 3 The application of the new coding system to describe double-rooted teeth with fused roots.

(Nusstein 2012, Luder 2015). Thus, the new system can be used to describe root fusions in two different ways:

1. Double-rooted teeth: If a researcher considers root fusion as a normal anatomical variation, then a slash (/) or double slash (//) can be added between the roots if the canals are separate or joined, respectively – without writing an abbreviation for root fusion on the left of the tooth number (Fig. 3). If a researcher considers root fusion
2. Three-rooted teeth: Root fusion in three-rooted teeth can have several presentations, as described by Zhang *et al.* (2014), which can be used in the new coding system as shown in Fig. 4. The use of one slash (/) is not required (Fig. 4a, b); the




Figure 4 Micro-CT images for the application of the new coding system to describe root fusion in three-rooted teeth. (a) A left maxillary first molar having fused MB and DB roots – Type 1 root fusion (MB fused to DB) according to Zhang *et al.* (2014) classification. RF – Root Fusion. (b) A maxillary molar tooth 18 with a Type 5 root fusion (MB and DB fused to P). (c) A maxillary tooth 28 with a type I root fusion in which the MB and DB canals are merged (double slash - //).

double slash (//) can be used if canals are merged (Fig. 4c).

When it is difficult to differentiate between a single root with deep developmental grooves, and roots that have complete fusion along the root length, then the categorization can be based on the common number of roots for that particular tooth. As an example, if a double-rooted maxillary premolar tooth has deep buccal and palatal grooves in the buccal root (Fig. 5), then it is considered as a buccal root with deep developmental grooves (not as two fused buccal roots) since it is well-known in dental anatomy that the most common root anatomy of maxillary first

premolars is either single or double rooted, in addition, the furcation groove in the palatal aspect of the buccal roots in double-rooted maxillary first premolars is considered as a normal anatomical landmark (Li *et al.* 2013). The same applies for maxillary lateral incisors and mandibular premolars with deep developmental grooves, which are considered as single-rooted, not fused double roots.

It is worth noting that classifications for deep developmental grooves have been described (Kogon 1986, Gu 2011), and they can also be applied in the new coding system as shown in Fig. 6. Other dental anomalies such as accessory roots (Fig. 7), and




Figure 5 Micro-CT reconstructed images showing double-rooted maxillary first premolar in which the buccal root has deep buccal and palatal grooves (B: Buccal, P: Palatal).




Figure 6 Micro-CT reconstructed images showing mandibular right first premolar with proximal-lingual groove type III (the groove is long (beyond the coronal third of the root) and deep, corresponding to a complex root canal system) (according to Gu 2011 classification; PGG – proximal-lingual groove).




Figure 7 Application of the new coding system to describe teeth with accessory roots. MB: Mesiobuccal, DB: Distobuccal, MP: Mesiopalatal, DP: Distopalatal.

anomalies that can occur in the furcation area such as enamel pearls, which can occur more than once on a tooth (Fig. 8), or along the root component such as root dilacerations (Fig. 9), as well as other rare root anomalies such as root fusions between teeth (concrescence – a cemental fusion of roots) can also be described using the new coding system (Fig. 10). The new coding system also allows researchers to describe the anomaly of interest, regardless of the presence of other anatomical variations if these are not related to the research (Fig. 11). The presence of different anomalies in the same tooth can also be described (Fig. 11).

On some occasions, a research study (related to anatomy or others related to mechanical instrumentation of roots having a single or double canals) will focus on a specific root such as the mesiobuccal or mesial root in maxillary or mandibular molar teeth,




Figure 8 Application of the new system to describe a tooth having more than one of the same anomaly. EP – Enamel Pearl.




Figure 9 The use of the new coding system to describe teeth with root dilacerations (RD). (a) Affecting one root (written beside that particular root) or (b) two roots (written before tooth number).




Figure 10 Micro-CT reconstructed images showing concrescence (C) between double-rooted mandibular left second molar with a supernumerary tooth (ST). M – Mesial, D – Distal. Note there is no communication between the pulp tissues of the two teeth.




Figure 11 The new coding system can classify teeth with more than one anatomical variation if it is an objective of the research. Micro-CT reconstructed image showing a maxillary molar tooth having two different anomalies – Enamel Pearl (EP) and root fusion (RF) type 4 (MB root fused to the DB and P roots) according to Zhang *et al.* (2014). [Colour figure can be viewed at wileyonlinelibrary.com]

respectively. In such cases, the codes can be written in a way that addresses this specification as shown in Fig. 12.

Root canal configuration

In order for a researcher to interpret the configuration of the root canal consistently, the components of the

pulp cavity, including the pulp chamber and root canal, should be defined accurately. This accurate interpretation has anatomical challenges because the transition from the pulp chamber to the root canal is not sharply demarcated macroscopically nor microscopically (Nelson & Ash 2010, Phulari 2019). According to the American Association of Endodontists (AAE 2020) and oral biology textbooks (Schied & Weiss




Figure 12 Examples for research studies which focus on (a) three-rooted maxillary first molars (MF), (b) only on sectioned MB roots of maxillary first molars, (c) only on MB roots having a single root canal regardless of the canal morphology of DB or P roots in maxillary first molars.




Figure 13 Micro-CT reconstructed images showing the level of the pulp tissue at CEJ (yellow line), and level of the furcation (floor of the pulp chamber; red line). In contrast to single-rooted teeth, the CEJ cannot be the level of the pulp chamber floor for double-/multi-rooted teeth.

2012, Nanci 2013), the pulp chamber is defined as the portion of the pulp space within the anatomic crown of the tooth. This could be a reasonable, consistent definition for single-rooted teeth, but it is not an accurate demarcation for double and multi-rooted teeth since the cemento enamel junction (CEJ) is not usually at the level of the floor of the pulp chamber (Nelson & Ash

2010, Nusstein 2012; Fig. 13), which is more often located some distance apical to the CEJ corresponding to the root trunk (Nelson & Ash 2010, Ahmed *et al.* 2018; Figs 13 and 14).

Most morphological studies (including common classifications for root and canal morphology) did not define where the floor of the pulp chamber ends, and




Figure 14 Micro-CT reconstructed images showing the location of the pulp chamber in both the crown and trunk of multi-rooted teeth.

where the orifice of the root canal begins to allow the canal configuration to be defined accurately. For instance, in Weine's classification (Weine *et al.* 1969), the floor of the pulp chamber was taken as a reference for the root canal orifice in the MB root of maxillary molars, but no information was given for the location of the orifice in single-rooted teeth. Similarly, Vertucci's classification did not define the location of the canal orifice, which was reported in a later publication (Vertucci 2005) as 'A root canal begins as a funnel-shaped canal orifice generally present *at or slightly apical to the cervical line*', with no clear definition of what 'slight apical' position actually means. One possible reason is that Vertucci's classification was based on the decalcification, staining and clearing method which may disrupt the normal anatomical features of the tooth significantly, including the CEJ, thus making its identification, in some samples, impossible.

The lack of a standard definition for the apical extent of the pulp chamber is a critical concern that can undermine the validity of comparisons amongst different studies which define root canal configurations with unclear, confusing and subjective anatomical landmarks. The new system for root and canal morphology defines the root canal configuration with a start (root canal orifice) passing through the canal and ends at the apical foramen. The root canal orifice is located at the level of the CEJ in single-rooted teeth, and at the level of root bi/trifurcation (floor of the pulp chamber) in double and multi-rooted teeth. On some occasions, the root bi/trifurcation in double-/multi-rooted teeth are located in the middle or apical third, in which a common canal is present coronally which starts from the level of CEJ, similar to single-rooted teeth. This common canal is written as a superscript before describing the canal configuration for each of the roots. Figures 15 and 16 show the




Figure 15 Micro-CT reconstructed images showing the general concept of the new coding system to define pulp chamber in single- and multi-rooted teeth.




Figure 16 Micro-CT reconstructed images showing the use of the new system to classify mandibular and maxillary teeth.

basic concepts of using the new coding system in different mandibular and maxillary permanent teeth.

Once the number of roots and location of root canal orifice(s) is defined, the root canal configuration can be interpreted systematically up to the foramen as described by Ahmed *et al.* (2017; Fig. 16); however, it has to be noted that the course of the canal (O-C-F) does not mean that each of the O, C and F should be allocated a number. The O-C-F system describes where the canal configuration starts and ends but the coding will be 1 if the root has a single canal throughout its length, 2 if the root has 2 canals, 1-2 if the configuration starts as 1 canal then

ends in 2 and so on, in a similar manner to previous classification systems (Weine *et al.* 1969, Vertucci 1984). However, the new system does not use Roman numerals and is therefore able to describe all canal systems, regardless of their complexity.

In micro-CT studies, it is common to find samples with a transverse canal anastomosis (isthmus), which is defined as a thin communication between two or more canals in the same root or between vascular elements in tissues (AAE 2020). Vertucci (1984) reported the percentage and location of a transverse canal anastomosis between canals as a separate entity from root canal configurations; however, the criteria




Figure 17 Micro-CT reconstruction of a right mandibular first premolar classified using the Vertucci classification with and without considering intercanal communication.

of defining such features were not mentioned, and the landmarks for differentiating it from canal configuration are unclear (Fig. 17). Others have classified the canal isthmus into subtypes based on characteristic morphological landmarks (Hsu & Kim 1997, von Arx 2005). Recent micro-CT studies have provided more detailed information on isthmuses (Keleş & Keskin 2018), and the ability of various irrigation/instrumentation protocols to remove hard tissue debris from these areas (Keleş *et al.* 2016). The confusion, nowadays, is more obvious when micro-CT studies continue to describe many canal configurations as 'nonclassifiable', which is true as a concept for some configurations but misleading and inappropriate for many other types because such studies have included transverse canals as a part of the configuration, which was not the case in the Vertucci classification (Fig. 17).

The new coding system defines transverse canal anastomosis as an integral part of the root canal configuration. The identification of transverse canal anastomosis separately from the canal configuration is a concern because they have clinical implications during chemo-mechanical instrumentation, canal filling and root-end cavity preparation and filling (von Arx 2005, Wu *et al.* 2006, Endal *et al.* 2011). In addition, transverse canal anastomosis may communicate with the external root surface via accessory canals and allow the transmission of toxins into the lateral periodontal and periapical tissues, thus affecting clinical outcomes (Villegas *et al.* 2004; Fig. 18). That is why the inclusion of intercanal communications as a part of the root canal configuration was considered in earlier classification systems of root canal morphology (Pineda 1973, Green 1973).

Despite the importance of adding intercanal communications to the root canal configuration, some




Figure 18 Micro-CT reconstructed image of a mandibular molar showing the connections of intercanal communications to the external root surface.




Figure 19 Micro-CT reconstructed image showing the use of the new coding system to describe teeth with complex intercanal communication. Asterisk indicates complex intercanal communication.

teeth may have ‘mesh-like’ transverse canals, or complex communications (Alavi *et al.* 2002), mainly caused by continuous deposition of secondary and

reparative dentine because of irritants and age changes, together with inherent genetic factors within different population groups; this will make classifying




Figure 20 Application of the new system to describe C-shaped canals (CSC). CSC¹ refers to C-shaped canals type I according to Fan *et al.* (2007) classification.

the root canal configuration of such teeth more complicated and probably impossible. To solve this, it is probably more appropriate to use an abbreviation or a code (such as an asterisk after describing the canal configuration or in-between the orifice and foramen to indicate a complex course of the canal configuration) when intercanal communications are complex and have multiple connections and branchings (Fig. 19). This will help researchers to allocate samples into specific groups in a more organized manner. The canal morphology of certain roots can also be tabulated separately such as MB^{O-C-F} and M^{O-C-F} for mesiobuccal and mesial roots in maxillary and mandibular molars, respectively. Other root canal anatomical aberrations can be described using the new system such as C-shaped canals (Fig. 20).

Similar to apical root bifurcations mentioned above, divisions of the main canal in the apical third are difficult to categorize. Common classifications for canal morphology did not provide clear definitions for such anatomy (Weine *et al.* 1969, Vertucci *et al.* 1974), and the assessment of apical canal configurations continues to vary amongst different observers, with some apical bifurcations classified either as an accessory canal or a clear division of the original single canal into two (type 1–2). Some may consider those canals deviating from the course of the main canal as an accessory, and some may consider accessory canals in the apical third as small bifurcating canals

that cannot be negotiated clinically and can only be observed after root filling (Ahmed *et al.* 2017). According to the AAE (2020), an accessory canal is defined as ‘any branch’ of the main pulp canal or chamber that communicates with the external surface of the root; this definition does not provide accurate characteristics for such canals and does not differentiate accessory canals from bifurcating main canals in the apical third of the root. It is obvious that, to date, a standard and consistent view of such anatomy has not yet been achieved (Ahmed *et al.* 2017).

Accessory canals

An accessory canal is a small canal leaving the main ‘root canal’ that (usually) communicates with the external surface of the root or furcation. As a consequence, it can be located anywhere along the length of the root (coronal, middle or apical third), or pulp chamber (chamber canals) and can be any type (patent, blind, loop; Fig. 21). An apical delta refers to multiple accessory canals (more than two) at or near the root apex (Ahmed *et al.* 2018; Fig. 21). The propagation of microbes and their by-products occurs not only within the main root canal but also in complex anatomical features that communicate with periradicular tissues, resulting in periodontitis anywhere along the length of the root, apex or furcation (Dammaschke *et al.* 2004, Ahmed 2012, Jang *et al.* 2015).




Figure 21 Types of accessory canals identified in micro-CT studies, and steps for the application of the new system to classify patent (configuration 1), blind (configuration 1-0) and loop canals (configuration 2-1-0) in addition to chamber canals in which the configuration is written between brackets.

The new coding system has complementary codes that can be added to the main code for canal configuration in order to provide details of the main and minor canal anatomy. There are two important landmarks for application in the new coding system to describe accessory canals – the location of the accessory canal (coronal, middle or apical third of the root)

and the number/configuration of the accessory canal [either patent (coded as 1), blind (coded as 1-0), loop (coded as 2-1-0) or apical delta with an abbreviation of 'D' (Ahmed *et al.* 2018)] as shown in Figs 21–23. Accessory canals are written after the canal configuration between brackets except for chamber canals which are written as a superscript before the roots




Figure 22 Using of the new coding system to describe accessory canals in micro-CT samples. C1 refers to the presence of one accessory canal (patent) in the coronal third of the root. A1 refers to the presence of one accessory canal (patent) in the apical third of the root. 1-0 refers to a blind accessory canal.




Figure 23 Application of the new coding system to describe accessory canals identified in more than one-third – M1, D refers to one accessory canal in the middle third and an apical delta (D). M refers to the mesial root.

(Fig. 21). Notably, accessory canals can occur in more than one-third, which also can be applied in the new system as shown in Fig. 23.

Application of the new coding system in the primary dentition

Root and canal morphology in the primary dentition is variable and complex (Ahmed 2013, 2014, Fumes *et al.* 2014, Ahmed *et al.* 2016, El Hachem *et al.* 2019). The new classification system is more accurate

compared with the Vertucci classification that was developed to categorize canal systems in permanent teeth (Ahmed *et al.* 2020b). Figure 24 shows the application of the new system to describe the root and canal morphology in the primary teeth scanned using micro-CT.

As mentioned above, the description of accessory canals using the new system in the permanent dentition is clear, and it can be applied in the primary dentition (Fig. 25). However, the observer may have challenges when interpreting canal morphology if the




Figure 24 Application of the new coding system to describe the root and canal morphology in anterior and posterior primary teeth. * refers to complex intercanal communications.

roots are at advanced stages of physiologic root resorption. Anomalies can be described as mentioned for the permanent dentition (Ahmed & Dummer 2018a, b; Fig. 26).

Step-by-step workflow of the new coding system in micro-CT studies

Figure 27 shows the workflow for using the new coding system in micro-CT studies starting from tooth

selection and progressing through the description of roots, main and accessory canal morphology together with dental anomalies.

Application of the new coding system in CBCT studies

Similar to micro-CT studies, the new coding system can be used in CBCT investigations – either in extracted teeth or *in vivo* (Saber *et al.* 2019, Karobari




Figure 25 Application of the new system to describe accessory canals in the primary dentition. M1 and M3 refer to the presence of one and three patent accessory canals in the middle third of the root, respectively.




Figure 26 Application of the new system to describe accessory roots in the primary dentition. RE^{II} refers to Radix Entomolaris type II (curvature in the coronal third and straight continuation to the apex) according to Song *et al.* (2010).

et al. 2020). For extracted teeth, the principles for applying the new system would follow the same as those for micro-CT studies mentioned above. It is well-known that, up to date, CBCT does not usually allow visualization of very fine details within canal systems (such as small branchings and accessory canals) compared to micro-CT; however, it is a valid diagnostic tool in clinical settings (Abella *et al.* 2015, Martins *et al.* 2017, Ahmed & Rossi-Fedele 2020).

In clinical CBCT studies, the number of roots (and abnormalities/anomalies such as root fusion, deep grooves, dilacerations and others) can be described. The identification of root canal orifices

and description of root canal configurations will follow the same principles as described in micro-CT studies (Fig. 28). The observer will have to determine the most apical position of the CEJ (cervical line) and level of pulp floor in single- and multi-rooted teeth, respectively, in order to determine the location of the orifice (Fig. 28). The presence of intercanal communications is usually not as complex as micro-CT studies; however, CBCT studies in different population groups have reported nonclassifiable types when they used the existing systems, but which can be classified using the new coding system (Karobari *et al.* 2020).




Figure 27 Step-by-step workflow of the new coding system.




Figure 28 The use of the new coding system to define the pulp chamber in CBCT scans.




Figure 29 Application of the new system to describe bifid root morphology in CBCT studies.

Because of their inherent limitations and the overlap of surrounding hard and soft tissues, fine details of the root such as the apex morphology (mainly the double apex morphology described above) may not be identified clearly in CBCT images. However, a bifid root apex can be identified as shown in Fig. 29. Accessory canals are not usually observed in CBCT studies; however, some accessory canals can be large

enough to be identified in CBCT images. The new coding system can be used to describe this minor anatomy (if identified) as shown in Fig. 30.

Figures 31–33 show the application of the new coding system in CBCT scans having maxillary and mandibular teeth. Figures 34 and 35 show the application of the new system to describe dental anomalies.




Figure 30 Describing the accessory canals (if identified) in CBCT studies using the new coding system.




Figure 31 Application of the new system to describe maxillary teeth in CBCT scans.




Figure 32 Application of the new system to describe mandibular teeth in CBCT scans.




Figure 33 Ability of the new system to classify nonclassifiable Vertucci canal configurations.

Apart from studies involving microCT and CBCT technology, the application of the new coding system in clinical practice (including different phases of clinical management from pre-operative

radiographic interpretation passing through intra-operative procedures to root canal filling) has recently been explained and elaborated (Ahmed et al. 2021).



Figure 34 Application of the new coding system to describe anomalies in the anterior dentition using CBCT. DI – Dens Invaginatus, DI^I and DI^{III} – Dens Invaginatus types I and III based on Oehlers (1957) classification; DE – Dens Evaginatus.




Figure 35 Application of the new coding system to describe other anomalies/variations in CBCT studies such as a) proximal gingival groove (PGG) type III, b) C-shaped canal (CSC) type II in which the M and D roots are fused (/) but each root has its own canal configuration and c) root fusion (RF) type 1 in which the MB and DB roots are fused and share the canal configuration (//) – starting as 3 canals and merging into 2 (3-2).

Conclusions

Despite variation in the accuracy of currently available technology and the clarity they bring to defining root and canal anatomy, the new coding system provides a clear description of the main and accessory canal morphology as well as anomalies in all tooth types when using micro-CT and CBCT. This article explains, elaborates and provides a detailed discussion on several controversial issues related to fine details of the root apex and canal bifurcations as well as anatomical landmarks of the root canal system that require a universal consensus for consistent presentation of root and canal anatomy.

Acknowledgements

Extracted teeth were collected and scanned using a micro-CT system (Xradia 520 Versa, Ziess, Germany) [Ethical approval number – DF RD2031/0140 (L)], Department of Geology, Faculty of Science, University of Malaya. 3D reconstruction was performed using Mimics software (Materialise, Leuven, Belgium). Images from CBCT machine (Kodak 9000 3D; Kodak Carestream Health, Trophy, France) were retrieved from the Department of Oral and Maxillofacial Clinical Sciences, Faculty of Dentistry, University of Malaya [Ethical approval number – DF RD1801/0004/1971/20152 (L)]. This project is supported by the following grants (Principal Investigator – HMA Ahmed): RU Grant number GPF018A-2020; BK Grant number BK-010-2018.

Conflict of interest

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

References

- AAE (2020) *Glossary of Terms*. American Association of Endodontists. <https://www.aae.org/specialty/download/glossary-of-endodontic-terms/>
- Abella F, Teixidó LM, Patel S, Sosa F, Duran-Sindreu F, Roig M (2015) Cone-beam computed tomography analysis of the root canal morphology of maxillary first and second premolars in a Spanish population. *Journal of Endodontics* **41**, 1241–7.
- Ahmed HMA (2012) Different perspectives in understanding the pulp and periodontal intercommunications with a new proposed classification for endo-perio lesions. *ENDO - Endodontic Practice Today* **6**, 87–104.
- Ahmed HMA (2013) Anatomical challenges, electronic working length determination and current developments in root canal preparation of primary molar teeth. *International Endodontic Journal* **46**, 1011–22.
- Ahmed HMA (2014) Pulpectomy procedures in primary molar teeth. *European Journal of General Dentistry* **3**, 3–10.
- Ahmed HMA, Abbott PV (2012) Accessory roots in maxillary molar teeth: a review and endodontic considerations. *Australian Dental Journal* **57**, 123–31; quiz 248.
- Ahmed HMA, Che Ab Aziz ZA, Azami NH et al. (2020a) Application of a new system for classifying root canal morphology in undergraduate teaching and clinical practice: a national survey in Malaysia. *International Endodontic Journal* **53**, 871–9.
- Ahmed HMA, Dummer PMH (2018a) A new system for classifying tooth, root and canal anomalies. *International Endodontic Journal* **51**, 389–404.
- Ahmed HMA, Dummer PMH (2018b) Advantages and applications of a new system for classifying roots and canal systems in research and clinical practice. *European Endodontic Journal* **3**, 9–17.
- Ahmed HMA, Hashem AAR, Dummer PMH (2021) Application of a new system for classifying root and canal anatomy in clinical practice – Explanation and elaboration. *European Endodontic Journal* In Press.
- Ahmed HMA, Khamis MF, Gutmann JL (2016) Seven root canals in a deciduous maxillary molar detected by the dental operating microscope and micro-computed tomography. *Scanning* **38**, 554–7.
- Ahmed HMA, Musale PK, El Shahawy OI, Dummer PMH (2020b) Application of a new system for classifying tooth, root and canal morphology in the primary dentition. *International Endodontic Journal* **53**, 27–35.
- Ahmed HMA, Neelakantan P, Dummer PMH (2018) A new system for classifying accessory canal morphology. *International Endodontic Journal* **51**, 164–76.
- Ahmed HMA, Rossi-Fedele G (2020) Preferred reporting items for root and canal anatomy in the human dentition (PROUD 2020) – a systematic review and a proposal for a standardized protocol. *European Endodontic Journal* **5**, 159–76.
- Ahmed HMA, Versiani MA, De-Deus G, Dummer PMH (2017) A new system for classifying root and root canal morphology. *International Endodontic Journal* **50**, 761–70.
- Alavi AM, Opananon A, Ng YL, Gulabivala K (2002) Root and canal morphology of Thai maxillary molars. *International Endodontic Journal* **35**, 478–85.
- von Arx T (2005) Frequency and type of canal isthmuses in first molars detected by endoscopic inspection during periradicular surgery. *International Endodontic Journal* **38**, 160–8.
- Badole GP, Warhadpande MM, Sheno PR, Lachure C, Badole SG (2014) A rare root canal configuration of bilateral maxillary first molar with 7 root canals diagnosed using cone-beam computed tomographic scanning: a case report. *Journal of Endodontics* **40**, 296–301.

- Dammaschke T, Witt M, Ott K, Schäfer E (2004) Scanning electron microscopic investigation of incidence, location, and size of accessory foramina in primary and permanent molars. *Quintessence International* **35**, 699–705.
- El Hachem C, Kaloustian MK, Nehme W *et al.* (2019) Three-dimensional modeling and measurements of root canal anatomy in second primary mandibular molars: a case series micro CT study. *European Archives of Paediatric Dentistry* **20**, 457–65.
- Endal U, Shen Y, Knut A, Gao Y, Haapasalo M (2011) A high-resolution computed tomographic study of changes in root canal isthmus area by instrumentation and root filling. *Journal of Endodontics* **37**, 223–7.
- Fan W, Fan B, Gutmann JL, Cheung GS (2007) Identification of C-shaped canal in mandibular second molars. Part I: radiographic and anatomical features revealed by intraradicular contrast medium. *Journal of Endodontics* **33**, 806–10.
- Filipo-Perez C, Bramante CM, Villas-Boas MH, Hungaro Duarte MA, Versiani MA, Ordinola-Zapata R (2015) Micro-computed tomographic analysis of the root canal morphology of the distal root of mandibular first molar. *Journal of Endodontics* **41**, 231–6.
- Fumes AC, Sousa-Neto MD, Leoni GB *et al.* (2014) Root canal morphology of primary molars: a micro-computed tomography study. *European Archives Paediatric Dentistry* **15**, 317–26.
- Green D (1973) Double canals in single roots. *Oral Surgery Oral Medicine Oral Pathology* **35**, 689–96.
- Gu YC (2011) A micro-computed tomographic analysis of maxillary lateral incisors with radicular grooves. *Journal of Endodontics* **37**, 789–92.
- Hashem AAR, Ahmed HMA (2017) Endodontic management of a mandibular first molar with unusual canal morphology. *European Endodontic Journal* **2**, 2–5.
- Hou GL, Tsai CC (1994) The morphology of root fusion in Chinese adults (I). Grades, types, location and distribution. *Journal of Clinical Periodontology* **21**, 260–4.
- Hsu YY, Kim S (1997) The resected root surface. The issue of canal isthmuses. *Dental Clinics of North America* **41**, 529–40.
- Jang JH, Lee JM, Yi JK, Choi SB, Park SH (2015) Surgical endodontic management of infected lateral canals of maxillary incisors. *Restorative Dentistry & Endodontics* **40**, 79–84.
- Kantiliaraki E, Delantoni A, Angelopoulos C, Beltes P (2019) Evaluation of root and root canal morphology of mandibular first and second molars in a Greek population: a CBCT study. *European Endodontic Journal* **4**, 62–8.
- Karobari MI, Noorani TY, Halim MS, Ahmed HMA (2020) Root and canal morphology of the anterior permanent dentition in Malaysian population using two classification systems: a CBCT clinical study. *Australian Endodontic Journal*. <https://doi.org/10.1111/aej.12454>
- Keleş A, Alçin H, Sousa-Neto MD, Versiani MA (2016) Supplementary steps for removing hard tissue debris from isthmus-containing canal systems. *Journal of Endodontics* **42**, 1677–82.
- Keleş A, Keskin C (2018) A micro-computed tomographic study of band-shaped root canal isthmuses, having their floor in the apical third of mesial roots of mandibular first molars. *International Endodontic Journal* **51**, 240–6.
- Keleş A, Keskin C, Alqawasm R, Versiani MA (2020) Micro-computed tomographic analysis of the mesial root of mandibular first molars with bifid apex. *Archives of Oral Biology* **117**, 104792.
- Kogon SL (1986) The prevalence, location and conformation of palato-radicular grooves in maxillary incisors. *Journal of Periodontology* **57**, 231–4.
- Kottoor J, Hemamalathi S, Sudha R, Velmurugan N (2010) Maxillary second molar with 5 roots and 5 canals evaluated using cone beam computerized tomography: a case report. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* **109**, e162–5.
- Leoni GB, Versiani MA, Pécora JD, Damião de Sousa-Neto M (2014) Micro-computed tomographic analysis of the root canal morphology of mandibular incisors. *Journal of Endodontics* **40**, 710–6.
- Li J, Li L, Pan Y (2013) Anatomic study of the buccal root with furcation groove and associated root canal shape in maxillary first premolars by using micro-computed tomography. *Journal of Endodontics* **39**, 65–8.
- Luder HU (2015) Malformations of the tooth root in humans. *Frontiers in Physiology* **6**, 307.
- Martins JNR, Marques D, Mata A, Caramês J (2017) Root and root canal morphology of the permanent dentition in a Caucasian population: a cone beam computed tomography study. *International Endodontic Journal* **50**, 1013–26.
- Nanci A (2013) *Ten Cate's Oral Histology*. St. Louis: Elsevier, pp. 165–204.
- Nelson SJ, Ash MM (2010) *Wheeler's Dental Anatomy, Physiology and Occlusion*. St. Louis: Elsevier, pp. 111–2.
- Nusstein JM (2012) Application of root and pulp morphology related to endodontic therapy. In: Scheid RC, Weiss G, eds. *Woelfel's Dental Anatomy*, 8th edn. Philadelphia: Lippincott Williams & Wilkins, pp. 231–49.
- Oehlers FA (1957) Dens invaginatus (dilated composite odontome). I. Variations of the invagination process and associated anterior crown forms. *Oral Surgery Oral Medicine Oral Pathology* **10**, 1204–18.
- Phulari RGS (2019) *Textbook of Dental Anatomy, Physiology and Occlusion*, 2nd edn. New Delhi: Jaypee Brothers Medical Publishers, pp. 238–48.
- Pineda F (1973) Roentgenographic investigation of the mesiobuccal root of the maxillary first molar. *Oral Surgery Oral Medicine Oral Pathology* **36**, 253–60.
- Ross IF, Evanchik PA (1981) Root fusion in molars: incidence and sex linkage. *Journal of Periodontology* **52**, 663–7.
- Saber SEDM, Ahmed MHM, Obeid M, Ahmed HMA (2019) Root and canal morphology of maxillary premolar teeth in an Egyptian subpopulation using two classification

- systems: a cone beam computed tomography study. *International Endodontic Journal* **52**, 267–78.
- Salas H, Torres J, Pauro J, Ahmed HMA (2021) A new coding system for root and canal morphology – an online survey from Peru. *International Endodontic Journal*, **54**, 147–9.
- Schied R, Weiss G (2012) *Woelfel's Dental Anatomy*, 8th edn. Philadelphia: Lippincott Williams & Wilkins, pp. 3–41.
- Song JS, Choi HJ, Jung IY, Jung HS, Kim SO (2010) The prevalence and morphologic classification of distolingual roots in the mandibular molars in a Korean population. *Journal of Endodontics* **36**, 653–7.
- Turner CG (1981) Root number determination in maxillary first premolars for modern human populations. *American Journal of Physical Anthropology* **54**, 59–62.
- Versiani MA, Ordinola-Zapata R, Keleş A et al. (2016) Middle mesial canals in mandibular first molars: a micro-CT study in different populations. *Archives of Oral Biology* **61**, 130–7.
- Versiani MA, Pécora JD, Sousa-Neto MD (2011) The anatomy of two-rooted mandibular canines determined using micro-computed tomography. *International Endodontic Journal* **44**, 682–7.
- Versiani MA, Pécora JD, de Sousa-Neto MD (2012) Root and root canal morphology of four-rooted maxillary second molars: a micro-computed tomography study. *Journal of Endodontics* **38**, 977–82.
- Vertucci FJ (1984) Root canal anatomy of the human permanent teeth. *Oral Surgery Oral Medicine Oral Pathology* **58**, 589–99.
- Vertucci F (2005) Root canal morphology and its relationship to endodontic procedures. *Endodontic Topics* **10**, 3–29.
- Vertucci F, Seelig A, Gillis R (1974) Root canal morphology of the human maxillary second premolar. *Oral Surgery Oral Medicine Oral Pathology* **38**, 456–64.
- Villegas JC, Yoshioka T, Kobayashi C, Suda H (2004) Frequency of transverse anastomoses with and without apical communication in Japanese population teeth. *Australian Endodontic Journal* **30**, 50–2.
- Weine FS, Hayami S, Hata G, Toda T (1999) Canal configuration of the mesiobuccal root of the maxillary first molar of a Japanese sub-population. *International Endodontic Journal* **32**, 79–87.
- Weine FS, Healey HJ, Gerstein H, Evanson L (1969) Canal configuration in the mesiobuccal root of the maxillary first molar and its endodontic significance. *Oral Surgery Oral Medicine Oral Pathology* **28**, 419–25.
- Wu MK, Dummer PMH, Wesselink PR (2006) Consequences of and strategies to deal with residual post-treatment root canal infection. *International Endodontic Journal* **39**, 343–56.
- Xu T, Fan W, Tay FR, Fan B (2019) Micro-computed tomographic evaluation of the prevalence, distribution, and morphologic features of accessory canals in Chinese permanent teeth. *Journal of Endodontics* **45**, 994–9.
- Xu T, Tay FR, Gutmann JL et al. (2016) Micro-computed tomography assessment of apical accessory canal morphologies. *Journal of Endodontics* **42**, 798–802.
- Zhang Q, Chen H, Fan B, Fan W, Gutmann JL (2014) Root and root canal morphology in maxillary second molar with fused root from a native Chinese population. *Journal of Endodontics* **40**, 871–5.